

Communication

# Strategic Recommendations to Mitigate Beirut Explosion Consequences

Ahmed Mohamed Abd El Mohimen Hasham <sup>1</sup>Department of Chemistry, Faculty of Science, Suez Canal University, Ismailia 41522, Egypt.

\*Correspondence: ahmed\_hesham@science.suez.edu.eg; Tel.: 00201159465989; ORCID: 0000-0002-0202-6664

**Abstract:** Beirut harbor explosion on 4th August 2020, carried out in the quarantine area of highly flammable substances caused, white gases emissions followed by reddish-brown gases emissions. The previous epidemiological studies stated that nitrogen dioxide has appeared to be a good indicator of the pollutant mixture. Furthermore, animal toxicological studies show that prolonged exposures can cause decreases in lung host defenses and changes in lung structure. Besides many adverse health effects on human organs like on blood, liver, and spleen. Also, cause hypertension, diabetes, heart, and cardiovascular diseases, and even death. Additionally, in July 2020, Ogen. Y stated that long-term exposure to NO<sub>2</sub> may be one of the most important contributors to fatality caused by the COVID-19 virus worldwide. On these grounds, it is proposed that a long-term recommendation for nitrogen dioxide be established. So, my recommendation to put a plan for monitoring the air quality parameters especially Nitrogen oxides, water quality parameters (drinking water, wastewater, and seawater), and soil parameters to develop mathematical models to draw the boundaries of the affected area and take the corrective actions accordingly.

**Keywords:** Beirut, explosion, environmental impact, hazardous, chemicals, strategic

**How to cite this paper:** Ahmed Mohamed Abd El Mohimen Hasham. Strategic Recommendations to Mitigate Beirut Explosion Consequences. *Multidisciplinary Journal of Health Sciences and Research* 2021, 1, 1-2. <https://doi.org/10.31586/health001>

Received: March 01, 2021

Accepted: April 20, 2021

Published: April 21, 2021



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The white gas emissions which were followed by reddish-brown gas emissions was big adequate to clip the attention of many people of Beirut residents who captured it on their cellphones. The last one was great and caused a shockwave and a sonic boom, cause many financial losses. Lebanese Prime Minister Hassan Diab supposed a probable 2750 tons of ammonium nitrate had been kept at a warehouse for six years ( [www.cnn.com](http://www.cnn.com) ) (CNN 2021). The direct impact it has killed more than 135 people and injured over 3000 in the Lebanese capital. Ammonium nitrate was behind many terror attacks like in Oklahoma City, US in 1995, also the same substance was behind a massive deadly blast in Tianjin, China in 2015.

## 2. Findings

The released toxic gases by the blast, urging people to stay indoors and wear masks with a filter to adsorb the gases on the filter media. The expected released gases from this explosion Nitrogen monoxide (NO), Nitrogen dioxide (NO<sub>2</sub>), and Dinitrogen tetroxide (N<sub>2</sub>O<sub>4</sub>). The nitrogen oxides will be predominately nitric oxide but this can gradually change to nitrogen dioxide as the effluent moves away from the fire [1].

Long-term exposure to NO<sub>2</sub> may be one of the most important contributors to fatality caused by the COVID-19 virus worldwide [2]. Also, it has been confirmed that the irregular variability of the diffusion of SARS-CoV-2 in Italy could partially depend on the levels of atmospheric pollutants. Even though chronic exposure to atmospheric pollutants and related diseases may represent a risk factor in determining the severity of COVID-19 syndrome and the high incidence of fatal events [3].

### 3. Environmental and Health Impact:

According to the US, Department of Labor, Occupational Safety, and Health Administration office, Nitrogen dioxide is categorized as a respiratory irritant and the route of exposure is mostly inhalation. The term "silo-fillers disease" is associated with exposure to NO<sub>2</sub> as well as other nitrogen oxides <https://www.osha.gov/> (Occupational Safety and Health Administration 2021)

There is only lonely indication of disorders of the mechanism of breathing and ventilatory role on repeated exposure to NO<sub>2</sub> concentrations of (1–5 ppm). The infinite common of lung biochemical studies show special effects only after acute or subchronic exposure to levels of nitrogen dioxide exceeding 2 ppm. The concentration (C) of nitrogen dioxide had more influence than exposure duration (T) [4].

While for long term exposure according to studies on animals had shown that several weeks to months of exposure to NO<sub>2</sub> concentrations of less than (1 ppm) cause a plethora of effects, primarily in the lung but also in other organs such as blood, liver, the spleen, hypertension, diabetes, heart and cardiovascular diseases and even death. Biochemical changes often reflect cellular alterations (lowest reported levels for several studies (0.2–0.4 ppm) but isolated cases of lower effective concentrations) (WHO, 2006) [5]. Nitrogen dioxide levels as low as (0.5 ppm) also increase defenselessness to bacterial and viral infection of the lung. While the expected concentration now in Beirut many thousands of times larger. The atmosphere became saturated by nitrogen oxides that may lead to acid rain and photochemical smog and will affect the soil, seawater especially those close to the Beirut harbor, drinking water, and wastewater stations.

More severe exposures (>50 ppm) are characterized by pulmonary edema, cyanosis, bronchiolitis obliterans, respiratory failure, and death. The LC<sub>50</sub> (Lethal Concentration 50) for a 4-hour exposure is approximately 90 ppm NO<sub>2</sub>. (Occupational Safety and Health Administration 2021). Some studies have stated that chronic exposure to NO<sub>2</sub> is linked with increased risk of all-cause death and proposed a specific association of the gas with cardiopulmonary mortality as well.

### 4. Obstacles

There is a high degree of scientific complications involved, very little research on the kinetics and metabolism of nitrogen dioxide has been conducted. The available information is inadequate and only partially describes its deposition and fate in the respiratory tract.

### 5. Recommendations

It is highly recommended to monitor the air, water, and soil parameters to develop a mathematical model to identify the affected area accurately. Also, it is recommended to make clinical studies for all persons who are exposed to the explosion in the near. COVID-19 case numbers to be followed in focus because it is expected to rise significantly.

### References

1. D. Fattorini and F. Regoli, "Role of the atmospheric pollution in the Covid-19 outbreak risk in Italy," *medRxiv*. 2020, doi: 10.1101/2020.04.23.20076455.
2. Y. Ogen, "Assessing nitrogen dioxide (NO<sub>2</sub>) levels as a contributing factor to coronavirus (COVID-19) fatality," *Sci. Total Environ.*, 2020, doi: 10.1016/j.scitotenv.2020.138605.
3. K. T. Paul, T. R. Hull, K. Lebek, and A. A. Stec, "Fire smoke toxicity: The effect of nitrogen oxides," *Fire Saf. J.*, 2008, doi: 10.1016/j.firesaf.2007.10.003.
4. R. Williams, P. Jones, C. Croghan, J. Thornburg, and C. Rodes, "The influence of human and environmental exposure factors on personal NO<sub>2</sub> exposures," *J. Expo. Sci. Environ. Epidemiol.*, 2012, doi: 10.1038/jes.2011.20.
5. World Health Organization, Air Quality Guidelines. Global update 2005. 2006.